

Proving the Triangle Congruence Theorems

Sentence Frames for Proofs

Transformations:

- Translate _____ from _____ to _____.
- Rotate _____ using _____ as the center by angle _____.
- Rotate _____ using _____ as the center so that _____ coincides with _____.
- Reflect _____ across _____.
- Reflect _____ across the perpendicular bisector of _____.
- Segments _____ and _____ are the same length so they are congruent. Therefore, there is a rigid motion that takes _____ to _____. Apply that rigid motion to _____.

Justifications:

- We know the image of _____ is congruent to _____ because rigid motions preserve measure.
- Points _____ and _____ coincide after translating because we defined our translation that way!
- Since points _____ and _____ are the same distance along the same ray from _____ they have to be in the same place.
- Rays _____ and _____ coincide after rotating because we defined our rotation that way!
- The image of _____ must be on ray _____ since both _____ and _____ are on the same side of _____ and make the same angle with it at _____.
- Points _____ and _____ coincide because they are both at the intersection of the same lines, and 2 distinct lines can only intersect in 1 place.
- _____ is the perpendicular bisector of the segment connecting _____ and _____, because the perpendicular bisector is determined by 2 points that are both equidistant from the endpoints of a segment.

Conclusion statement:

- We have shown that a rigid motion takes _____ to _____, _____ to _____, and _____ to _____, therefore triangle _____ is congruent to triangle _____.

What Do We Know For Sure About Isosceles Triangles?

Kiran

Kiran: I'm stumped on this proof.

Mai: What are you trying to prove?

Kiran: I'm trying to prove that in an isosceles triangle, the two base angles are congruent. So in this case, that angle A is congruent to angle B .

Mai: Let's think of what geometry ideas we already know are true.

Kiran: We know if two pairs of corresponding sides, and the corresponding angles between the sides, are congruent, then the triangles must be congruent.

Mai: Yes, and we also know that we can use reflections, rotations, and translations to prove congruence and symmetry. . . The isosceles triangle you've drawn makes me think of symmetry. If you draw a line down the middle of it, I wonder if that could help us prove that the angles are the same?

[Mai draws the line of symmetry of the triangle and labels the intersection of AB and the line of symmetry Q .]

Kiran: Wait, when you draw the line, it breaks the triangle into two smaller triangles. I wonder if I could prove those triangles are congruent using Side-Angle-Side Congruence?

Mai: It's an isosceles triangle, so we know that one pair of corresponding sides is congruent. [Mai marks the congruent sides.]

Kiran: And this segment in the middle here is part of both triangles, so it has to be the same length for both. Look.

[**Kiran draws** the two halves of the isosceles triangle and marks the shared sides as congruent.]

Mai: So we have two pairs of corresponding sides that are congruent. How do we know the angles between them are congruent?

Kiran: I'm not sure. Maybe it has to do with how we drew that line of symmetry?

What Do We Know For Sure About Isosceles Triangles?

Mai

Kiran: I'm stumped on this proof.

Mai: What are you trying to prove?

Kiran: I'm trying to prove that in an isosceles triangle, the two base angles are congruent. So in this case, that angle A is congruent to angle B .

Mai: Let's think of what geometry ideas we already know are true.

Kiran: We know if two pairs of corresponding sides, and the corresponding angles between the sides, are congruent, then the triangles must be congruent.

Mai: Yes, and we also know that we can use reflections, rotations, and translations to prove congruence and symmetry. . . The isosceles triangle you've drawn makes me think of symmetry. If you draw a line down the middle of it, I wonder if that could help us prove that the angles are the same?

[**Mai draws** the line of symmetry of the triangle and labels the intersection of AB and the line of symmetry Q .]

Kiran: Wait, when you draw the line, it breaks the triangle into two smaller triangles. I wonder if I could prove those triangles are congruent using Side-Angle-Side Congruence?

Mai: It's an isosceles triangle, so we know that one pair of corresponding sides is congruent.

[Mai marks the congruent sides.]

Kiran: And this segment in the middle here is part of both triangles, so it has to be the same length for both. Look.

[Kiran draws the two halves of the isosceles triangle and marks the shared sides as congruent.]

Mai: So we have two pairs of corresponding sides that are congruent. How do we know the angles between them are congruent?

Kiran: I'm not sure. Maybe it has to do with how we drew that line of symmetry?