

# Solving Exponential Equations

Let's solve equations using logarithms.

## 14.1 A Valid Solution?

Here is a solution to the equation  $5 \cdot e^{3a} = 90$ .

$$\begin{aligned} 5 \cdot e^{3a} &= 90 \\ e^{3a} &= 18 \\ 3a &= \log_e 18 \\ a &= \frac{\log_e 18}{3} \end{aligned}$$

Explain what happened in each step.

## 14.2 Natural Logarithm

1. Complete the table with equivalent equations. The first row is completed for you.

	exponential form	logarithmic form
	$e^6 \approx 403.43$	$\ln(403.43) \approx 6$
a.	$e^0 = 1$	
b.	$e^1 = e$	
c.	$e^{-1} = \frac{1}{e}$	
d.		$\ln \frac{1}{e^2} = -2$
e.	$e^x = 10$	

2. Solve each equation by expressing the solution using  $\ln$  notation. Then, find the approximate value of the solution using the “ $\ln$ ” button on a calculator.

a.  $e^m = 20$

b.  $e^n = 30$

c.  $e^p = 7.5$

## 14.3 Solving Exponential Equations

Without using a calculator, solve each equation. It is expected that some solutions will be expressed using log notation. Be prepared to explain your reasoning.

$$1. \ 10^x = 10,000$$

$$2. \ 5 \cdot 10^x = 500$$

$$3. \ 10^{(x+3)} = 10,000$$

$$4. \ 10^{2x} = 10,000$$

$$5. \ 10^x = 315$$

$$6. \ 2 \cdot 10^x = 800$$

$$7. \ 10^{(1.2x)} = 4,000$$

$$8. \ 7 \cdot 10^{(0.5x)} = 70$$

$$9. \ 2 \cdot e^x = 16$$

$$10. \ 10 \cdot e^{3x} = 250$$



 **Are you ready for more?**

Assume that  $a$  and  $b$  are positive values. Use your understanding of what logarithms mean to find these values. Explain or show your reasoning.

1.  $\log_a(a^b)$

2.  $a^{\log_a(b)}$

## Lesson 14 Summary

So far we have solved exponential equations by

- Finding whole number powers of the base (for example, the solution to  $10^x = 100$  is  $x = 2$ , and the solution to  $10^y = 1,000$  is  $y = 3$ ).
- Estimation (for example, the solution of  $10^x = 300$  is between 2 and 3 because 300 is between 100 and 1,000).
- Using a logarithm and approximating its value on a calculator (for example, the solution of  $10^x = 300$  is  $\log 300 \approx 2.48$ ).

Sometimes solving exponential equations takes additional reasoning. Here are a couple of examples.

$$\begin{array}{ll} 5 \cdot 10^x = 45 & 10^{(0.2t)} = 1,000 \\ 10^x = 9 & 10^{(0.2t)} = 10^3 \\ x = \log 9 & 0.2t = 3 \\ & t = \frac{3}{0.2} \\ & t = 15 \end{array}$$

In the first example, the power of 10 is multiplied by 5, so to find the value of  $x$  that makes this equation true, each side is divided by 5. From there, the equation is rewritten as a logarithm, giving an exact value for  $x$ .

In the second example, the expressions on each side of the equation are rewritten as powers of 10:  $10^{(0.2t)} = 10^3$ . This means that the exponent  $0.2t$  on one side and the 3 on the other side must be equal, and leads to an expression to solve where we don't need to use a logarithm.

How do we solve an exponential equation with base  $e$ , such as  $e^x = 5$ ? We can express the solution using the **natural logarithm**, the logarithm for base  $e$ . Natural logarithm is written as  $\ln$ , or sometimes as  $\log_e$ . Just like the equation  $10^2 = 100$  can be rewritten, in logarithmic form, as  $\log_{10} 100 = 2$  or  $\log 100 = 2$ , the equation  $e^0 = 1$  can be rewritten as  $\ln 1 = 0$ . Similarly,  $e^{-2} = \frac{1}{e^2}$  can be rewritten as  $\ln \frac{1}{e^2} = -2$ .

All this means that we can solve  $e^x = 5$  by rewriting the equation as  $x = \ln 5$ . This says that  $x$  is the exponent to which base  $e$  is raised to equal 5.

To estimate the size of  $\ln 5$ , remember that  $e$  is about 2.7. Because 5 is greater than  $e^1$ , this means that  $\ln 5$  is greater than 1.  $e^2$  is about  $(2.7)^2$ , or 7.3. Because 5 is less than  $e^2$ , this means that  $\ln 5$  is less than 2. This suggests that  $\ln 5$  is between 1 and 2. Using a calculator we can check that  $\ln 5 \approx 1.61$ .