Different Options for Solving One Equation

Let's think about which way is better when we solve equations with parentheses.

9.1 Math Talk: Solve Each Equation

Solve each equation mentally.

- $100(x - 3) = 1{,}000$
- $500(x - 3) = 5{,}000$
- $0.03(x - 3) = 0.3$
- $0.72(x + 2) = 7.2$

 CC BY NC Illustrative Mathematics, based on IM 6–8 Math, CC BY Open Up Resources.

9.2 Analyzing Solution Methods

Three students each attempted to solve the equation $2(x-9)=10$, but they got different solutions. Here is their work. Do you agree with any of their methods? Explain or show your reasoning.

Noah's method:

$$\begin{aligned} 2(x-9) &= 10 \\ 2(x-9)+9 &= 10+9 && \text{Add 9 to each side} \\ 2x &= 19 \\ 2x \div 2 &= 19 \div 2 && \text{Divide each side by 2} \\ x &= \frac{19}{2} \end{aligned}$$

Elena's method:

$$\begin{aligned} 2(x-9) &= 10 \\ 2x-18 &= 10 && \text{Apply the distributive property} \\ 2x-18-18 &= 10-18 && \text{Subtract 18 from each side} \\ 2x &= -8 \\ 2x \div 2 &= -8 \div 2 && \text{Divide each side by 2} \\ x &= -4 \end{aligned}$$

Andre's method:

$$\begin{aligned} 2(x-9) &= 10 \\ 2x-18 &= 10 && \text{Apply the distributive property} \\ 2x-18+18 &= 10+18 && \text{Add 18 to each side} \\ 2x &= 28 \\ 2x \div 2 &= 28 \div 2 && \text{Divide each side by 2} \\ x &= 14 \end{aligned}$$

CC BY NC Illustrative Mathematics, based on IM 6–8 Math, CC BY Open Up Resources.

9.3 Solution Pathways

1. Solve each of these equations twice, one time using each method.

   a. applying the distributive property first:

   $$2{,}000(x - 0.03) = 6{,}000$$

   dividing each side first:

   $$2{,}000(x - 0.03) = 6{,}000$$

   b. applying the distributive property first:

   $$2(x + 1.25) = 3.5$$

   dividing each side first:

   $$2(x + 1.25) = 3.5$$

 CC BY NC Illustrative Mathematics, based on IM 6–8 Math, CC BY Open Up Resources.

2. Solve each of these equations once. Choose whichever method you think will be easier for that equation.

   a. $\frac{1}{4}(4 + x) = \frac{4}{3}$

   b. $-10(x - 1.7) = -3$

   c. $5.4 = 0.3(x + 8)$

 CC BY NC Illustrative Mathematics, based on IM 6–8 Math, CC BY Open Up Resources.

Lesson 9 Summary

Equations can be solved in many ways. In this lesson, we focused on equations with a specific structure, and two specific ways to solve them.

Suppose we are trying to solve the equation $\frac{4}{5}(x+27)=16$. Two useful approaches are:

- Divide each side by $\frac{4}{5}$.
- Apply the distributive property.

In order to decide which approach is better, we can look at the numbers and think about which would be easier to compute. We notice that $\frac{4}{5} \cdot 27$ will be hard, because 27 isn't divisible by 5. So, distributing the $\frac{4}{5}$ is not the best method. But $16 \div \frac{4}{5}$ gives us $16 \cdot \frac{5}{4}$, and 16 is divisible by 4. So, dividing each side by $\frac{4}{5}$ is a good choice.

$$\begin{aligned} \frac{4}{5}(x+27) &= 16 \\ \frac{5}{4} \cdot \frac{4}{5}(x+27) &= 16 \cdot \frac{5}{4} \\ x+27 &= 20 \\ x &= -7 \end{aligned}$$

Sometimes the calculations are simpler if we first use the distributive property. Let's look at the equation $100(x+0.06)=21$. If we first divide each side by 100, we get $\frac{21}{100}$ or 0.21 on the right side of the equation. But if we use the distributive property first, we get an equation that only contains whole numbers.

$$\begin{aligned} 100(x+0.06) &= 21 \\ 100x+6 &= 21 \\ 100x &= 15 \\ x &= \frac{15}{100} \end{aligned}$$

 CC BY NC Illustrative Mathematics, based on IM 6–8 Math, CC BY Open Up Resources.