


Lesson 16 Practice Problems

1. There are many cylinders with a volume of 144π cubic inches. The height $h(r)$ in inches of one of these cylinders is a function of its radius r in inches where $h(r) = \frac{144}{r^2}$.
 - a. What is the height of one of these cylinders if its radius is 2 inches?
 - b. What is the height of one of these cylinders if its radius is 3 inches?
 - c. What is the height of one of these cylinders if its radius is 6 inches?


2. The surface area $S(r)$ in square units of a cylinder with a volume of 18 cubic units is a function of its radius r in units where $S(r) = 2\pi r^2 + \frac{36}{r}$. What is the surface area of a cylinder with a volume of 18 cubic units and a radius of 3 units?

3. Han finds an expression for $S(r)$ that gives the surface area in square inches of any cylindrical can with a specific fixed volume, in terms of its radius r in inches. This is the graph Han gets if he allows r to take on any value between -1 and 5.

- a. What would be a more appropriate domain for Han to use instead?
- b. What is the approximate minimum surface area for the can?



4. The graph of a polynomial function f is shown. Is the degree of the polynomial even or odd? Explain your reasoning.



(From Unit 2, Lesson 8.)

5. The polynomial function $p(x) = x^4 + 4x^3 - 7x^2 - 22x + 24$ has known factors of $(x + 4)$ and $(x - 1)$.
- Rewrite $p(x)$ as the product of linear factors.
 - Draw a rough sketch of the graph of the function.

(From Unit 2, Lesson 12.)

6. Which polynomial has $(x + 1)$ as a factor?

A. $x^3 + 2x^2 - 19x - 20$

B. $x^3 - 21x + 20$

C. $x^3 + 8x + 11x - 20$

D. $x^3 - 3x^2 + 3x - 1$

(From Unit 2, Lesson 15.)